

GENTAMED- gentamicin sulfate powder
Bimeda, Inc.

DESCRIPTION: Each gram of GentaMed™ Soluble Powder contains gentamicin sulfate equivalent to 333.33 mg gentamicin base.

RESIDUE WARNING: For use in swine drinking water only. Swine treated with the recommended dose of GentaMed™ Soluble Powder must not be slaughtered for food for at least ten (10) days following the last treatment.

INDICATIONS: GentaMed™ Soluble Powder is recommended for the control and treatment of colibacillosis in weanling swine caused by strains of *E. coli* sensitive to gentamicin and for the control and treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

DOSAGE AND ADMINISTRATION:

GentaMed Soluble Powder is available in a 360 gram jar. The 360 gram jar contains a scoop that provides approximately 18 grams of product when level full.

For *Control and treatment of Colibacillosis*-Administer GentaMed Soluble Powder in drinking water at the recommended level of 25 mg per gallon (1 scoop/240 gallons) for three consecutive days.

For *Control and treatment of Swine Dysentery*-Administer GentaMed Soluble Powder in drinking water at the recommended level of 50 mg per gallon (1 scoop/120 gallons) for three consecutive days.

The concentration of the medication should be adjusted in extremely hot or cold weather to ensure a gentamicin dosage of approximately 0.5 mg/lb/day for 3 days for colibacillosis or 1.0 mg/lb/day for 3 days for swine dysentery. A 50-lb pig will, under average conditions, consume 0.9 gallons of water per day.

For swine dysentery, if the condition recurs, the medication may be repeated. Because of the tendency for the disease to recur on a premise with a history of swine dysentery, a control program should be used following treatment.

DIRECTIONS FOR USE:

To prepare medicated water, mix GentaMed Soluble Powder and drinking water according to the following tables:

Proportioner Use:

For Control and Treatment of Colibacillosis

Add 18 g (1 scoop) GentaMed Soluble Powder to 2 gallons of water to make stock solution and dispense at the rate of one ounce per gallon of drinking water. This will result in drinking water that contains 25 mg/gallon.

For Control and Treatment of Swine Dysentery

Add 18 g (1 scoop) GentaMed Soluble Powder to 1 gallon of water to make stock solution and dispense at the rate of one ounce per gallon of drinking water. This will result in drinking water that contains 50 mg/gallon.

Bulk Preparation:

For Control and Treatment of Colibacillosis

Add 18 g (1 scoop) GentaMed Soluble Powder to 240 gallons of water.

For Control and Treatment of Swine Dysentery

Add 18 g (1 scoop) GentaMed Soluble powder to 120 gallons of water.

Medicated drinking water should not be stored or offered in rusty containers since the drug is quickly destroyed in such containers.

Medicated water should be prepared daily and be the sole source of drinking water for 3 consecutive days.

The daily water consumption figures shown below are approximations for mild temperatures and are presented as a guide only. Actual water consumption varies with environmental temperature, humidity, and diet.

Approximate Daily Water Consumption

Weight of Pig (lb)	Gallon for 5 Pigs	Gallons for 10 Pigs
25	3.0	6.0
50	4.5	9.0
75	6.0	12.0
100	7.5	15.0

SIDE EFFECTS: There have been no reported side effects following the recommended use of GentaMed™ Soluble Powder in swine.

CONTRAINDICATIONS: There are no known contraindications to this drug when used as directed.

PRECAUTIONS: Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).

HOW SUPPLIED: GentaMed™ Soluble Powder is available in a 360 gram jar with 2 jars per shipper.

GentaMed™ Soluble Powder

(Gentamicin Sulfate)

Contains 120 grams of Gentamicin per jar

Antibacterial

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Restricted Drug (California)-Use Only as Directed

ANADA 200-494, Approved by FDA

Net weight: 360 g (12.7 oz)

GentaMed™ Soluble Powder (Gentamicin Sulfate) ANTIBACTERIAL

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
FOR SWINE DRINKING WATER ONLY

DESCRIPTION: Each gram of GentaMed™ Soluble Powder contains gentamicin sulfate equivalent to 333.33 mg gentamicin base.

RESIDUE WARNING: For use in swine drinking water only. Swine treated with the recommended dose of GentaMed™ Soluble Powder must not be slaughtered for food for at least ten (10) days following the last treatment.

INDICATIONS: GentaMed™ Soluble Powder is recommended for the control and treatment of colibacillosis in weanling swine caused by strains of *E. coli* sensitive to gentamicin and for the control and treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

DOSE AND ADMINISTRATION: GentaMed™ Soluble Powder is available in a 360 gram jar. The 360 gram jar contains a scoop that provides approximately 18 grams of product when level full.

For Control and treatment of Colibacillosis-Administer GentaMed™ Soluble Powder in drinking water at the recommended level of 25 mg per gallon (1 scoop/240 gallons) for three consecutive days.

For Control and treatment of Swine Dysentery- Administer GentaMed™

Soluble Powder in drinking water at the recommended level of 50 mg per gallon (1 scoop/120 gallons) for three consecutive days.

The concentration of the medication should be adjusted in extremely hot or cold weather to ensure a gentamicin dosage of approximately 0.5 mg/lb/day for 3 days for colibacillosis or 1.0 mg/lb/day for 3 days for swine dysentery. A 50-lb pig will, under average conditions, consume 0.9 gallons of water per day.

For swine dysentery, if the condition recurs, the medication may be repeated. Because of the tendency for the disease to recur on a premise with a history of swine dysentery, a control program should be used following treatment.

DIRECTIONS FOR USE: To prepare medicated water, mix GentaMed™ Soluble Powder and drinking water according to the following tables:

Proportioner Use

For Control and Treatment of Colibacillosis
Add 18 g (1 scoop) GentaMed™ Soluble Powder to 2 gallons of water to make stock solution and dispense at the rate of one ounce per gallon of drinking water. This will result in drinking water that contains 25 mg/gallon.

For Control and Treatment of Swine Dysentery
Add 18 g (1 scoop) GentaMed™ Soluble Powder to 1 gallon of water to make stock solution and dispense at the rate of

one ounce per gallon of drinking water. This will result in drinking water that contains 50 mg/gallon.

Bulk Preparation:

For Control and Treatment of Colibacillosis
Add 18 g (1 scoop) GentaMed™ Soluble Powder to 240 gallons of water.

For Control and Treatment of Swine Dysentery
Add 18 g (1 scoop) GentaMed™ Soluble Powder to 120 gallons of water.

Medicated drinking water should not be stored or offered in rusty containers since the drug is quickly destroyed in such containers.

Medicated water should be prepared daily and be the sole source of drinking water for 3 consecutive days.

The daily water consumption figures shown below are approximations for mild temperatures and are presented as a guide only. Actual water consumption varies with environmental temperature, humidity, and diet.

Approximate Daily Water Consumption

Weight of Pig (lb)	Gallons for 5 Pigs	Gallons for 10 Pigs
25	3.0	6.0
50	4.5	9.0
75	6.0	12.0
100	7.5	15.0

SIDE EFFECTS: There have been no reported side effects following the recommended use of GentaMed™ Soluble Powder in swine.

CONTRAINDICATIONS: There are no known contraindications to this drug when used as directed.

PRECAUTIONS: Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C-30°C (59°F-86°F).

HOW SUPPLIED: GentaMed™ Soluble Powder is available in a 360 gram jar with 2 jars per shipper.

For Animal Use Only

Keep Out of Reach of Children
Restricted Drug (California)-Use Only as Directed

EACH g CONTAINS: Gentamicin sulfate equivalent to 333.33 mg gentamicin base.

RESIDUE WARNING: Swine treated with the recommended dose of GentaMed™ Soluble Powder must not be slaughtered for food for at least ten (10) days following the last treatment.

Do not store or offer medicated water in rusty metal containers.

Read accompanying directions carefully. Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C-30°C (59°F-86°F).

For use in swine drinking water only.
8GEN001 Rev. 05/16 Product No. 1GEN009



Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com



INDICATIONS: GentaMed™ Soluble Powder is recommended for the control and treatment of colibacillosis in weanling swine caused by strains of *E. coli* sensitive to gentamicin, and for the control and treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

DOSE:

Colibacillosis – 1 level scoop per 240 gallons of drinking water for 3 consecutive days, 0.5 mg/lb/day. For proportioner use, add 1 level scoop to 2 gallons stock solution and dispense at the rate of 1 oz/gal drinking water (25 mg/gal).

Swine Dysentery – 1 level scoop per 120 gallons of drinking water for 3 consecutive days, 1.0 mg/lb/day. For proportioner use, add 1 level scoop to 1 gallon stock solution and dispense at the rate of 1 oz/gal drinking water (50 mg/gal).

GentaMed is a Trademark of Bimeda, Inc.

EACH g CONTAINS: Gentamicin sulfate equivalent to 333.33 mg gentamicin base.

RESIDUE WARNING: Swine treated with the recommended dose of GentaMed™ Soluble Powder must not be slaughtered for food for at least ten (10) days following the last treatment.

Do not store or offer medicated water in rusty metal containers.

Read accompanying directions carefully. Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C-30°C (59°F-86°F).

For use in swine drinking water only.
8GEN001 Rev. 05/16 Product No. 1GEN009



PULL HERE TO OPEN

TAKE TIME
OBSERVE LABEL DIRECTIONS

GENTAMED

gentamicin sulfate powder

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:6 1133-5135
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
gentamicin sulfate (UNII: 8X7386QRLV) (GENTAMICIN - UNII:T6Z9V48IKG)	GENTAMICIN	120 g in 360 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:6 1133-5135-1	360 g in 1 JAR		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200494	12/14/2011	

Labeler - Bimeda, Inc. (060492923)

Registrant - Bimeda, Inc. (060492923)

Establishment

Name	Address	ID/FEI	Business Operations
Bimeda, Inc.		060492923	manufacture

Revised: 11/2018

Bimeda, Inc.