

Joint Support for Dogs

TriCOX™



TECHNICAL SPECIFICATIONS: Joint care formula containing UC-II, astaxanthin, glucosamine, MSM and creatine in a palatable liver flavored soft chew.

ACTIVE INGREDIENTS (per 1 chew):

- Glucosamine HCL (marine source):450 mg
- MSM (Mehtylsulfonylmethane):400 mg
- Creatine Monohydrate:250 mg
- Bromelain:75 mg
- Boswellia serrata extract:20 mg
- UC-II®:10 mg
- Eicosapentaenoic Acid (EPA):9 mg
- Docosahexaenoic Acid (DHA):6 mg
- Astaxanthin 3 mg

INACTIVE INGREDIENTS: Black Malt Extract, Caramel Color, Carob Powder, Citric Acid, Dry Brewer's Yeast, Dry Poultry Liver Digest, Dried Chicken Liver, Fish Oil, Gluconolactone, Glycerin, Hydrolyzed Wheat Protein, Maltodextrine, Mixed Tocopherols (preservative), Natural Flavor, Poultry Fat, Rosemary Extract, Papain, Sodium Metabiosulfite (preservative), Sorbic Acid (preservative), Soy Flour, Soy Lecithin, Stevia, Sweet Whey, Tapioca Starch, Wheat Flour, Wheat Starch.

PROPERTIES:

TriCOX™ contains a unique blend of ingredients to help support the joint function in many ways. UC-II is a patented form of undenatured collagen that helps maintain a normal immune response to prevent auto-immune attacks on the joint.

Boswellia extract helps maintain normal inflammatory response within the joint.

Astaxanthin is a powerful antioxidant that has been proven to be 6,000 times more potent than vitamin C.

TriCOX™ also brings the cartilage building blocks glucosamine and MSM, and muscle support with creatine.

TriCOX™ is formulated using the TRP Soft Chew technology, giving the product unparalleled softness and palatability.

SPECIES: Dogs

DIRECTIONS FOR USE:

- 5 to 15 lbs: ½ chew every other day
 - 16 to 30 lbs: ½ chew daily
 - 31 to 50 lbs: 1 chew daily
 - 51 to 100 lbs: 2 chews daily
 - Over 100 lbs: 3 chews daily
- Dosage should be doubled during the initial period of 4 to 6 weeks

CAUTION: Safe use in pregnant animal or animals intended for breeding has not been proven. Consult with your veterinarian before using this product in animals with clotting disorders, being treated with anticoagulant medications; diabetes, or any other metabolic disorder causing hyperglycemia; history or urinary tract stones; known allergy to shellfish. If lameness worsens, discontinue use and contact your veterinarian. Administer during meal or after the animal has eaten to reduce incidence of gastrointestinal upset. An examination from a veterinarian is recommended prior to using this product.

STORAGE: Store at controlled room temperature.

WARNINGS: For animal use only. Keep out of reach of children and animals. In case of accidental overdose, contact a health professional immediately.

PRESENTATION: 60ct and 120ct tub.

UC-II is a registered trademark of InterHealth N.I.



Veterinarians can purchase VetOne® products exclusively through MWI Veterinary Supply or AAHA MARKETLink.

