



zoetis

Lincomix®

brand of lincomycin injection



WARNING

Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment. **Not for human use.**

CAUTION

If no improvement is noted within 48 hours, consult a veterinarian.

ADVERSE REACTIONS

The intramuscular administration to swine may cause a transient diarrhea or loose stools. Although this effect has rarely been reported, one must be alert to the possibility that it may occur.

Should this occur, it is important that the necessary steps be taken to prevent the effects of dehydration.

DOSAGE AND ADMINISTRATION

For arthritis or mycoplasma pneumonia—5 mg per pound of body weight intramuscularly once daily for three to seven days as needed. When using LINCOMIX Injectable containing 100 mg/mL, 1 mL/20 lb body weight will provide 5 mg/lb. When using LINCOMIX Injectable containing 300 mg/mL, 1 mL/60 lb body weight will provide 5 mg/lb.

For optimal results, initiate treatment as soon as possible.

As with any multi-dose vial, practice aseptic techniques in withdrawing each dose. Adequately clean and disinfect the vial closure prior to entry with a sterile needle and syringe. No vial closure should be entered more than 20 times.

INDICATIONS FOR SWINE

LINCOMIX Injectable is indicated for the treatment of infectious forms of arthritis caused by organisms sensitive to its activity. This includes most of the organisms responsible for the various infectious arthritides in swine, such as the staphylococci, streptococci, *Erysipelothrix* and *Mycoplasma spp.*

It is also indicated for the treatment of mycoplasma pneumonia.

CONTRAINDICATIONS

As with all drugs, the use of LINCOMIX Injectable is contraindicated in animals previously found to be hypersensitive to the drug.



HOW SUPPLIED

LINCOMIX Injectable is available in two concentrations: 300 mg/mL and 100 mg/mL.

300 mg/mL: *For use in swine weighing 300 pounds or more.*

Each mL contains lincomycin hydrochloride equivalent to lincomycin, 300 mg; also Benzyl Alcohol, 9.45 mg added as preservative. Supplied in 100 mL vials.

100 mg/mL: Each mL contains lincomycin hydrochloride equivalent to lincomycin, 100 mg; also Benzyl Alcohol, 9.45 mg added as preservative. Supplied in 100 mL vials.

Store at controlled room temperature 20°- 25°C (68°- 77°F), with excursions between 15°- 40°C (59°- 104°F).

Use contents within 28 days of first vial broach.

Approved by FDA under NADA # 034-025

zoetis

Distributed by:

Zoetis Inc.

Kalamazoo, MI 49007

Revised: May 2019