

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com

Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

activyl® PROTECTOR BAND FOR DOGS

Intervet/Merck Animal Health

ADJUSTABLE ONE-SIZE-FITS ALL

FLEA & TICK COLLAR FOR DOGS

- **Patented insecticide-release technology**
- **Kills fleas & ticks for up to 6 months**
- **Water Resistant**

Not affected by water

Active Ingredient:	Percentage by Weight
Deltamethrin	4.0%
Inert Ingredients	96.0%
Total	100.0%

UP TO 6 MONTHS PROTECTION

CAUTION

Do Not Let Children Play With This Collar

READ ENTIRE LABEL BEFORE EACH USE

- **USE ONLY ON DOGS**

Activyl Protector Band for Dogs provides full season, up to 6 months protection against fleas and ticks

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION:

Do not open protective pouch until ready to use. Do not let children play with this collar. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Do not use on puppies under 12 weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated, or nursing animals. Sensitivities may occur after using ANY pesticide products for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. This collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Applying other pesticides on the dog may not be necessary while the collar is being worn.

FIRST AID

IF SWALLOWED

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 - 20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

For incidents involving **HUMANS**, call 1-800-680-9206. For questions or incidents involving **ANIMALS**, call 1-800-224-5318 between the hours of 8am and 5pm. Evenings and weekends call 800-345-4735. Have the product container or label with you when calling a poison control center or doctor, or going for treatment

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **Wipe the collar with a damp cloth prior to application.** Place the collar around dog's neck, buckle and adjust for proper fit. Cut off approximately 2 inches from the buckle and dispose of excess length by wrapping in newspaper and placing in trash. The collar must be worn loosely so that two fingers may be placed between collar and dog's neck. Maximum effectiveness may not occur for 2-3 weeks after collar placement. Living and rest areas of pet must also be treated with appropriate pest control measures to ensure control of pests. Wetting will not impair the collar's effectiveness or the pet's protection. If the dog goes swimming or is out in the rain, it is not necessary to remove the collar. Activyl Protector Band for Dogs may be used in addition to a lead or constraint collar. Use only one Activyl Protector Band for Dogs at a time. This collar should be worn continuously. Reapply a new collar every 6 months.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store this product in a cool, dry place away from children and pets.

Pesticide and Container Disposal: Do not reuse container or used collar. Dispose of in trash.

IMPORTANT NOTICE: DISCLAIMER

Read "IMPORTANT NOTICE: DISCLAIMER" before buying or using. If terms are not acceptable, return at once unopened. To the extent consistent with applicable law, INTERVET INC. warrants only that the product conforms to the chemical description on the label and is reasonably fit for the purpose stated on the label when used in the accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to INTERVET INC., and user assumes the risk of any such use. To the extent consistent with applicable law, INTERVET INC. MAKES NO OTHER WARRANTY, EXPRESSED OR IMPLIED, INCLUDING ANY IMPLIED WARRANTY OR FITNESS FOR A PARTICULAR PURPOSE OR OF MERCHANTABILITY. In no case shall INTERVET INC. be liable for consequential, special, indirect or incidental damages resulting from the use of handling of this product. The foregoing conditions of sale and warranty can be varied only by an agreement in writing signed by a duly authorized representative of INTERVET INC.

Product of France

EPA Reg. No. 68451-1

EPA Est. no. 68451-FRA-1

Distributed by: Intervet Inc, d/b/a Merck Animal Health), 2 Giralda Farms, Madison, NJ 07940

NET CONTENTS:	NET WT.	
1 Collar	0.9 oz	179309 R1

CPN: 1047538.0

By using this content, you agree to the [Terms of Service](#) and [Privacy Policy](#).

Copyright 2021 - Animalytix LLC